5

144

## CLAIMS

1. Polyhydroxyalkanoate comprised of at least a unit represented by a chemical formula (1) within the molecule:

$$\begin{array}{c}
R\\N-H\\
=O\\
(CH_2)m\\
-\left(-O-\frac{|I|}{I}\right)-\\
Z
\end{array}$$
(1)

wherein R represents -A<sub>1</sub>-SO<sub>2</sub>R<sub>1</sub>; R<sub>1</sub> represents OH, a halogen atom, ONa, OK or OR<sub>1a</sub>; R<sub>1a</sub> and A<sub>1</sub> each independently represents a group having a substituted or unsubstituted aliphatic hydrocarbon structure, a substituted or unsubstituted aromatic ring structure or a substituted or unsubstituted heterocyclic structure; m represents an integer selected from 0 - 8; Z represents a linear or branched alkyl group, an aryl group or an aralkyl group substituted with an aryl group; and in case plural units are present, R, R<sub>1</sub>, R<sub>1a</sub>, A<sub>1</sub>, m and Z have the aforementioned meanings independently for each unit.

Polyhydroxyalkanoate according to claim 1,
 comprised of, as the unit represented by the chemical formula (1), at least a unit represented by a chemical formula (2), a chemical formula (3), a chemical formula (4A) or (4B), within a molecule:

145

$$\begin{array}{c} SO_2R_2 \\ A_2 \\ N-H \\ C=O \\ (CH_2)m \\ O \end{array}$$

wherein R<sub>2</sub> represents OH, a halogen atom, ONa, OK or OR<sub>2a</sub>; R<sub>2a</sub> represents a linear or branched alkyl group with 1 to 8 carbon atoms or a substituted or unsubstituted phenyl group; A<sub>2</sub> represents a linear or branched alkylene group with 1 to 8 carbon atoms; m represents an integer selected from 0 - 8; Z<sub>2</sub> represents a linear or branched alkyl group, an aryl group or an aralkyl group substituted with an aryl group; and in case plural units are present, A<sub>2</sub>, R<sub>2</sub>, R<sub>2a</sub>, m and Z<sub>2</sub> have the aforementioned meanings independently for each unit;

$$R_{3b}$$
 $R_{3c}$ 
 $R_{3d}$ 
 $R_{3e}$ 
 $R_{3e}$ 

wherein  $R_{3a}$ ,  $R_{3b}$ ,  $R_{3c}$ ,  $R_{3d}$  and  $R_{3e}$  each independently represents  $SO_2R_{3f}$  ( $R_{3f}$  representing OH, a halogen atom, ONa, OK or  $OR_{3f1}$  ( $R_{3f1}$  representing a linear or branched

PCT/JP2005/011000

alkyl group with 1 to 8 carbon atoms or a substituted or unsubstituted phenyl group)), a hydrogen atom, a halogen atom, an alkyl group with 1 - 20 carbon atoms, an alkoxy group with 1 - 20 carbon atoms, an OH group, an NH<sub>2</sub> group, an NO<sub>2</sub> group, COOR<sub>3g</sub> (R<sub>3g</sub> representing a H atom, a Na atom or a K atom), an acetamide group, an OPh group, a NHPh group, a CF<sub>3</sub> group, a C<sub>2</sub>F<sub>5</sub> group or a C<sub>3</sub>F<sub>7</sub> group (Ph indicating a phenyl group), of which at least one is SO<sub>2</sub>R<sub>3f</sub>; m represents an integer selected from 0 - 8; Z<sub>3</sub> represents a linear or branched alkyl group, an aryl group or an aralkyl group substituted with an aryl group; and in case plural units are present, R<sub>3a</sub>, R<sub>3b</sub>, R<sub>3c</sub>, R<sub>3d</sub>, R<sub>3e</sub>, R<sub>3f</sub>, R<sub>3f1</sub>, R<sub>3g</sub>, m and Z<sub>3</sub> have the aforementioned meanings independently for each unit;

$$\begin{array}{c|c} R_{4e} \\ R_{4f} \\ R_{4g} \\ R_{4a} \\ R_{4b} \\ R_{4d} \\ R_{$$

5

10

15

20

wherein  $R_{4a}$ ,  $R_{4b}$ ,  $R_{4c}$ ,  $R_{4d}$ ,  $R_{4e}$ ,  $R_{4f}$  and  $R_{4g}$  each independently represents  $SO_2R_{4o}$  ( $R_{4o}$  representing OH, a halogen atom, ONa, OK or  $OR_{4o1}$  ( $R_{4o1}$  representing a linear or branched alkyl group with 1 to 8 carbon

atoms or a substituted or unsubstituted phenyl group)), a hydrogen atom, a halogen atom, an alkyl group with 1 - 20 carbon atoms, an alkoxy group with 1 - 20 carbon atoms, an OH group, an  $\mathrm{NH}_2$  group, an  $\mathrm{NO}_2$ 5 group,  $COOR_{4p}$  ( $R_{4p}$  representing a H atom, a Na atom or a K atom), an acetamide group, an OPh group, an NHPh group, a  $CF_3$  group, a  $C_2F_5$  group or a  $C_3F_7$  group (Ph indicating a phenyl group), of which at least one is  $SO_2R_{4o}$ ; m represents an integer selected from 0 - 8; 10  $Z_{4a}$  represents a linear or branched alkyl group, an aryl group or an aralkyl group substituted with an aryl group; and in case plural units are present, R4a,  $R_{4b}$ ,  $R_{4c}$ ,  $R_{4d}$ ,  $R_{4e}$ ,  $R_{4f}$ ,  $R_{4g}$ ,  $R_{4o}$ ,  $R_{4o1}$ ,  $R_{4p}$ , m and  $Z_{4a}$  have the aforementioned meanings independently for each 15 unit;

20

wherein  $R_{4h}$ ,  $R_{4i}$ ,  $R_{4j}$ ,  $R_{4k}$ ,  $R_{4l}$ ,  $R_{4m}$  and  $R_{4n}$  each independently represents  $SO_2R_{4o}$  ( $R_{4o}$  representing OH, a halogen atom, ONa, OK or  $OR_{4o1}$  ( $R_{4o1}$  representing a linear or branched alkyl group with 1 to 8 carbon atoms or a substituted or unsubstituted phenyl

group)), a hydrogen atom, a halogen atom, an alkyl group with 1 - 20 carbon atoms, an alkoxy group with 1 - 20 carbon atoms, an OH group, an NH<sub>2</sub> group, an NO<sub>2</sub> group, COOR<sub>4p</sub> (R<sub>4p</sub> representing a H atom, a Na atom or a K atom), an acetamide group, an OPh group, an NHPh group, a CF<sub>3</sub> group, a C<sub>2</sub>F<sub>5</sub> group or a C<sub>3</sub>F<sub>7</sub> group (Ph indicating a phenyl group), of which at least one is SO<sub>2</sub>R<sub>4o</sub>; m represents an integer selected from 0 - 8; Z<sub>4b</sub> represents a linear or branched alkyl group, an aryl group or an aralkyl group substituted with an aryl group; and in case plural units are present, R<sub>4h</sub>, R<sub>4i</sub>, R<sub>4j</sub>, R<sub>4k</sub>, R<sub>4l</sub>, R<sub>4m</sub>, R<sub>4n</sub>, R<sub>4o</sub>, R<sub>4o1</sub>, R<sub>4p</sub>, m and Z<sub>4b</sub> have the aforementioned meanings independently for each unit.

3. Polyhydroxyalkanoate comprised of at least a unit represented by a chemical formula (5) within a molecule:

$$\begin{array}{c}
COOR_5 \\
(CH_2)m \\
O \\
-CO \\
Z_5
\end{array}$$
(4)

wherein R<sub>5</sub> represents hydrogen, a group capable of forming a salt or R<sub>5a</sub>; R<sub>5a</sub> represents a linear or branched alkyl group with 1 - 12 carbon atoms, an aralkyl group or a substituent having a sugar; m represents an integer selected from 0 - 8; Z<sub>5</sub> represents a linear or branched alkyl group, an aryl

WO 2005/121208

5

10

25

149

PCT/JP2005/011000

group or an aralkyl group substituted with an aryl group; however  $R_5$  only represents a substituent having a sugar in case  $Z_5$  is a methyl group and m is 0-1; and in case plural units are present,  $R_5$ ,  $R_{5a}$ , m and  $Z_5$  have the aforementioned meanings independently for each unit.

4. Polyhydroxyalkanoate according to any one of claims 1 to 3, further comprised of a unit represented by a chemical formula (6) within a molecule:

wherein R<sub>6</sub> represents a linear or branched alkylene with 1 - 11 carbon atoms, alkyleneoxyalkylene group (each alkylene group being independently with 1 - 2 carbon atoms), a linear or branched alkenyl group with 1 - 11 carbon atoms or an alkylidene group with 1 - 5 carbon atoms which may be substituted with an aryl group; and in case plural units are present, R<sub>6</sub> has the aforementioned meanings independently for each unit.

5. A method for producing a polyhydroxyalkanoate comprising a unit represented by a chemical formula (8), comprised of a step of executing hydrolysis of a polyhydroxyalkanoate comprising a unit represented by a chemical formula

WO 2005/121208

5

150

PCT/JP2005/011000

(7) in the presence of an acid or an alkali, or a step of executing hydrogenolysis comprising a catalytic reduction of a polyhydroxyalkanoate comprising a unit represented by a chemical formula (7):

$$\begin{array}{c}
COOR_7 \\
(CH_2)m \\
O \\
-
\end{array}$$

$$\begin{array}{c}
Z_7
\end{array}$$
(7)

wherein R<sub>7</sub> represents a linear or branched alkyl group with 1 - 12 carbon atoms or an aralkyl group; m represents an integer selected from 0 - 8; Z<sub>7</sub>

10 represents a linear or branched alkyl group, an aryl group or an aralkyl group substituted with an aryl group, and m represents an integer selected from 2 - 8 in case Z<sub>7</sub> is a methyl group; and in case plural units are present, R<sub>7</sub>, m and Z<sub>7</sub> have the

15 aforementioned meanings independently for each unit;

$$\begin{array}{c|c}
COOR_8 \\
(CH_2)m \\
O & \downarrow \\
Z_8
\end{array}$$
(8)

wherein R<sub>8</sub> represents hydrogen, or a group capable of forming a salt; m represents an integer selected from 0 - 8; Z<sub>8</sub> represents a linear or branched alkyl group, 20 an aryl group or an aralkyl group substituted with an aryl group, and m represents an integer selected from

2-8 in case  $Z_8$  is a methyl group; and, in case plural units are present,  $R_8$ , m and  $Z_8$  have the aforementioned meanings independently for each unit.

6. A method for producing a

5 polyhydroxyalkanoate comprising a unit represented by a chemical formula (1), comprised of a step of executing a condensation reaction of a polyhydroxyalkanoate comprising a unit represented by a chemical formula (9) and an amine compound 10 represented by a chemical formula (10):

$$\begin{array}{c}
\mathsf{COOR_9} \\
(\mathsf{CH_2})\mathsf{m} \\
\mathsf{O} \\
-\left(-\mathsf{O} \begin{array}{c} | & | \\ | & | \\ Z_{\mathsf{g}} \\ \end{array}\right) \\
(9)$$

wherein R<sub>9</sub> represents hydrogen, or a group capable of forming a salt; m represents an integer selected from 0 - 8; Z<sub>9</sub> represents a linear or branched alkyl group, an aryl group or an aralkyl group substituted with an aryl group; and, in case plural units are present, m, R<sub>9</sub> and Z<sub>9</sub> have the aforementioned meanings independently for each unit;

$$20 H_2N - A_3 - SO_2R_{10} (10)$$

wherein  $R_{10}$  represents OH, a halogen atom, ONa, OK or  $OR_{10a}$ ;  $R_{10a}$  and  $A_3$  each independently is selected from a group having a substituted or unsubstituted aliphatic hydrocarbon structure, a substituted or unsubstituted

152

aromatic ring structure, or a substituted or unsubstituted heterocyclic structure; and, in case plural units are present,  $R_{10}$ ,  $R_{10a}$  and  $A_3$  have the aforementioned meanings independently for each unit;

$$\begin{array}{c}
R\\N-H\\ = O\\ (CH_2)m\\ - O \end{array}$$

$$\begin{array}{c}
CH_2 = 0\\ CH$$

5

20

wherein R represents -A<sub>1</sub>-SO<sub>2</sub>R<sub>1</sub>; R<sub>1</sub> represents OH, a halogen atom, ONa, OK or OR<sub>1a</sub>; R<sub>1a</sub> and A<sub>1</sub> each independently represents a group having a substituted or unsubstituted aliphatic hydrocarbon structure, a substituted or unsubstituted aromatic ring structure or a substituted or unsubstituted heterocyclic structure; m represents an integer selected from 0 - 8; Z represents a linear or branched alkyl group, an aryl group or an aralkyl group substituted with an aryl group; and in case plural units are present, R, R<sub>1</sub>, R<sub>1a</sub>, A<sub>1</sub>, m and Z have the aforementioned meanings independently for each unit.

7. A method for producing a polyhydroxyalkanoate comprising a unit represented by a chemical formula (13), comprised of:

a step of reacting a polyhydroxyalkanoate comprising a unit represented by a chemical formula (11) with a base; and

153

a step of reacting a compound obtained in the aforementioned step with a compound represented by a chemical formula (12):

$$\begin{pmatrix} 0 & 1 & 1 \\ Z_{11} & 0 & 1 \end{pmatrix}$$
 (11)

wherein  $Z_{11}$  represents a linear or branched alkyl group, an aryl group or an aralkyl group substituted with an aryl group; and in case plural units are present,  $Z_{11}$  has the aforementioned meanings independently for each unit;

10

15

$$X(CH_2)mCOOR_{12}$$
 (12)

wherein m represents an integer selected from 0-8; X represents a halogen atom; and  $R_{12}$  represents a linear or branched alkyl group with 1-12 carbon atoms or an aralkyl group;

$$\begin{array}{c}
COOR_{13} \\
(CH_2)m \\
O \\
\downarrow II \\
Z_{13}
\end{array}$$
(13)

wherein m represents an integer selected from 0 - 8;

R<sub>13</sub> represents a linear or branched alkyl group with 1
- 12 carbon atoms or an aralkyl group; Z<sub>13</sub> represents

20 a linear or branched alkyl group, an aryl group or an aralkyl group substituted with an aryl group, and m represents an integer selected from 2 - 8 in case Z<sub>13</sub>

is a methyl group; and in case plural units are present,  $R_{13}$ , m and  $Z_{13}$  have the aforementioned meanings independently for each unit.

- 8. A method for producing a
- 5 polyhydroxyalkanoate comprising a unit represented by a chemical formula (15), comprised of:

a step of reacting a polyhydroxyalkanoate comprising a unit represented by a chemical formula (11) with a base; and

a step of reacting a compound obtained in the aforementioned step with a compound represented by a chemical formula (14):

$$-\left(-0\right)$$

wherein  $Z_{11}$  represents a linear or branched alkyl group, an aryl group or an aralkyl group substituted with an aryl group; and in case plural units are present,  $Z_{11}$  has the aforementioned meanings independently for each unit;

15

Ċ

wherein  $R_{14}$  represents  $-A_{14}-SO_2R_{14a}$ ;  $R_{14a}$  represents OH, a halogen atom, ONa, OK or  $OR_{14b}$ ;  $R_{14b}$  and  $A_{14}$  each independently is selected from a group having a substituted or unsubstituted aliphatic hydrocarbon

155

structure, a substituted or unsubstituted aromatic ring structure or a substituted or unsubstituted heterocyclic structure; and in case plural units are present,  $R_{14}$ ,  $R_{14a}$ ,  $R_{14b}$ , and  $A_{14}$  have the

5 aforementioned meanings independently for each unit;

$$\begin{array}{c}
R_{15} \\
N-H \\
= 0 \\
(CH_2)_{20} \\
-\left(-0 \right| 1 \\
Z_{15}
\end{array}$$
(15)

wherein R<sub>15</sub> represents -A<sub>15</sub>-SO<sub>2</sub>R<sub>15a</sub>; R<sub>15a</sub> represents OH, a halogen atom, ONa, OK or OR<sub>15b</sub>; R<sub>15b</sub> and A<sub>15</sub> each independently represents a group having a substituted or unsubstituted aliphatic hydrocarbon structure, a substituted or unsubstituted aromatic ring structure or a substituted or unsubstituted heterocyclic structure; Z<sub>15</sub> represents a linear or branched alkyl group, an aryl group or an aralkyl group substituted with an aryl group; and in case plural units are present, R<sub>15</sub>, R<sub>15a</sub>, R<sub>15b</sub>, and A<sub>15</sub> have the aforementioned meanings independently for each unit.